

High Q^2 HERA Events and pQCD at High x

Stephen Rock and Peter Bosted

American University, Washington D.C. 20016

June 1997

Abstract

We compare data on F_2^p from SLAC experiments in the range $0.7 \leq x \leq 0.97$ in and near the resonance region with previous empirical fits to Deep Inelastic Scattering and with calculations from parton distribution functions. The data is in rough agreement with the empirical fits, but is an order of magnitude higher than the pQCD calculations at the highest x . This compares with the two orders of magnitude increase in the quark distributions at high x which seem to be necessary to explain the HERA high Q^2 events.

Recent interest in the anomalous high Q^2 , high x events at HERA [1] has generated interest in the accuracy of the pQCD evolution to these regions. This evolution depends on $\int_x^1$ of the quark distributions, thus making the high x quark distributions important. pQCD fits have ignored data in the region $x \geq 0.75$ because of fear of possible higher twist contamination. However, there is a wealth of SLAC data in the region up to $x = 0.97$ [2, 3], albeit in the resonance region ($W^2 \leq 4\text{GeV}^2$) and at relatively low Q^2 . Bloom-Gilman [4] duality states that the average of the data in the resonance region approximates the DIS structure. With that in mind, we compare pQCD and various fits to the DIS region with the SLAC high- x resonance data.

Figure 1 shows a sample of the SLAC measurements of F_2^p for $x \geq 0.7$ and $7 \leq Q^2 \leq 30$. F_2^p decreases by approximately 3 orders of magnitude over this x range and there is a Q^2 dependence of approximately a factor of 2 to 4 between the various bands of data.

Figure 2 shows the ratio the same SLAC data to the NMC DIS fit [5]. This fit used only data with $x \leq 0.75$. The data in Figure 2. span the range $7 \leq Q^2 \leq 30$. In the range of validity of the fit ($x \leq 0.75$), the ratio is similar to unity, as expected. The ratio raises to about 2, at the highest x . This fit is dominated by a $(1-x)^3$ behavior at high x as predicted by the quark counting rules [6]. The agreement is remarkable considering that the data at the highest x is in the region of the first resonance, far from the normal

DIS region. The SLAC global fits used data up to $x=0.85$. As seen in Figure 3, similar good agreement is obtained when using the SLAC- Λ_{12} [3] as the denominator. This fit is also dominated by a $(1-x)^3$ behavior at high x . As observed in Figure 4, the ratio of data to the SLAC- Ω_9 fit is close to unity for the high Q^2 data, but the ratio is less than unity for the lower Q^2 , high- x points. Thus previous fits to world DIS data can be used to estimate the F_2^p structure function in the resonance region to within a factor of about two.

Figure 5 shows the ratio of SLAC data to a typical pQCD evolution fit, CTEQ 4M [7]. Again, there is good agreement for $x \approx 0.7$. However, the ratio raises sharply and is about a factor of 10 at the highest x . Note that since the data shown in Figure 1 decreases by a factor of 1000 over this range in x , this indicates that the pQCD fits decrease by 4 orders of magnitude. The difference between the pQCD fit and the data could be explained by “higher twist” effects, but they would have to be an order of magnitude greater than the pQCD evolution and also behave similar to the quark counting rule prediction of $(1-x)^3$. A typical higher twist form is $F_2^{pQCD}[1 + C_{HT}/(Q^2(1-x))]$ [8]. Figure 6 shows the ratio of the data to this higher twist form with $C_{HT} = 4$. While the rapid rise at high x from pQCD alone is considerably suppressed, the DIS region of excellent pQCD fits ($x \approx 0.7$) is suppressed by an unacceptably large 30%. A previous higher twist analysis [9] for $x \leq 0.75$ determined that the equivalent of C_{HT} is ≈ 0.3 , an order of magnitude smaller than that used in Figure 6.. Of course, more complex higher twist terms could be included with higher powers of $Q^2(1-x)$ which would eventually fit the data.

Recently Kuhlmann, Lai and Tung [10] have tried to explain the HERA events using a pQCD toy model which enhances the u quark probability beyond the CTEQ4M fit by approximately two orders of magnitude at $x=0.95$. This pQCD fit would overestimate the SLAC data by an order of magnitude and thus require a huge negative “higher twist effect” of approximately the form discussed above.

In conclusion, the SLAC data in the kinematic region $.75 \leq x \leq .97$ which is mostly in the resonance region, is an order of magnitude higher than current pQCD parton distributions, and an order of magnitude less than the toy models created to explain the high Q^2 HERA events. Existing DIS fits with quark counting rule behavior at high x are a reasonable approximation of the data, consistent with Bloom-Gilman duality. A dedicated experiment with the SLAC 50 GeV electron beam could measure F_2^p in the kinematic region $0.80 \leq x \leq 0.95$ in the region of $Q^2 \geq 50\text{GeV}^2$

with the possibility of tagging charm. This data could be used to disentangle higher twist effects from pQCD evolution, and determine the u and c quark distributions at high x .

References

- [1] H1 Collaboration (C. Adloff *et. al.*, DESY-97-024, Feb 1997. ZEUS Collaboration (J.Breitweg *et. al.*, DESY-97-025, Feb. 1997
- [2] P. E. Bosted *et. al.*, Phys. Rev. **D49**, 3091 (1994). S. Rock *et. al.*, Phys. Rev. **D46**, 24 (1992).
- [3] L. W. Whitlow *et. al.*, Phys. Lett. **bf B282**, 475 (1992).
- [4] E. D. Bloom and F. J. Gilman, Phys. Rev. Lett. **25**, 1140 (1970).
- [5] M. Arneodo *et. al.*, Phys. Lett. **B364**, 107 (1995).
- [6] S. J. Brodsky and M. Burkardt, Nucl. Phys. **B441**,197 (1995).
- [7] CERN Parton Distribution Library.
- [8] Richard Blankenbecler, Stanley J. Brodsky, Phys. Rev. **D10**,2973(1974)
- [9] Marc Virchaux, Alain Milsztajn, Phys.Lett. **B274**,221 (1992).
- [10] S. Kuhlmann, H. L. Lai, and W. K. Tung, MSU-HEP-70316, CTEQ-705, hep-ph/9704338 v2.




Figure 1: Selected SLAC data for F_2^p in the range $0.7 \leq x \leq 0.97$ and $7 \leq Q^2 \leq 30$. Most of the data at the higher x values is in the resonance region.

FIGURE 2: F_2 DATA/ F_2 -NMC




Figure 2: The data of Figure 1 divided by the NMC fit to Deep Inelastic Scattering Data for $x \leq 0.75$.

FIGURE 3: F_2 DATA/ $F_2 - \Lambda_{12}$




Figure 3: The data of Figure 1 for F_2^p divided by the SLAC Global Fit Λ_{12} to SLAC Deep Inelastic Scattering Data for $x \leq 0.85$.




Figure 4: The data of Figure 1 for F_2^p divided by the SLAC Global Fit Ω_9 to SLAC Deep Inelastic Scattering Data for $x \leq 0.85$.

FIGURE 5 F_2 DATA/ F_2 -pQCD-CTEQ 4M




Figure 5: The data of Figure 1 for F_2^p divided by the pQCD NLO fit 4M done by CTEQ.

FIGURE 6: F_2 DATA/ F_2 -pQCD-CTEQ 4M +HT=4.0




Figure 6: The data of Figure 1 for F_2^p divided by the pQCD NLO fit 4M done by CTEQ with an additional empirical higher twist term described in the text.